

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 August 2002 (01.08.2002)

PCT

(10) International Publication Number
WO 02/058702 A1

(51) International Patent Classification⁷: **A61K 31/47**,
31/4709, C07D 215/38, 401/12, 405/12, A61P 9/00

(21) International Application Number: PCT/US02/02007

(22) International Filing Date: 25 January 2002 (25.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/264,439 26 January 2001 (26.01.2001) US

(71) Applicant (for all designated States except US):
SMITHKLINE BEECHAM CORPORATION
[GB/GB]; One Franklin Plaza, Philadelphia, PA 19103
(GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DHANAK,**
Dashyant [GB/US]; 709 Swedeland Road, King of
Prussia, PA 19406 (US). **KNIGHT, Steven, D.** [US/US];
709 Swedeland Road, King of Prussia, PA 19406 (US).
WARREN, Gregory, L. [US/US]; 1250 South Col-
legeville Road, Collegeville, PA 19426 (US).

(74) Agents: **HALL, Linda, E.** et al.; SmithKline Beecham
Corporation, Corporate Intellectual Property, UW2220,
709 Swedeland Road, P.O. Box 1539, King of Prussia, PA
19406-0939 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: UROTENSIN-II RECEPTOR ANTAGONISTS

(57) Abstract: The present invention relates to quinolines, pharmaceutical compositions containing them and their use as antagonists of urotensin II.

WO 02/058702 A1

UROTENSIN-II RECEPTOR ANTAGONISTS

FIELD OF THE INVENTION

- 5 The present invention relates generally to quinolines, pharmaceutical compositions containing them, and their use as antagonists of urotensin II.

BACKGROUND OF THE INVENTION

- 10 The integrated control of cardiovascular homeostasis is achieved through a combination of both direct neuronal control and systemic neurohormonal activation. Although the resultant release of both contractile and relaxant factors is normally under stringent regulation, an aberration in this *status quo* can result in cardiohemodynamic dysfunction with pathological consequences.

- 15 The principal mammalian vasoactive factors that comprise this neurohumoral axis, namely angiotensin-II, endothelin-1, norepinephrine, all function via an interaction with specific G-protein coupled receptors (GPCR). Urotensin-II, represents a novel member of this neurohumoral axis.

In the fish, this peptide has significant hemodynamic and endocrine actions in diverse end-organ systems and tissues:

- 20 • smooth muscle contraction
both vascular and non-vascular in origin including smooth muscle preparations from the gastrointestinal tract and genitourinary tract. Both pressor and depressor activity has been described upon systemic administration of exogenous peptide
- 25 • osmoregulation:
effects which include the modulation of transepithelial ion (Na^+ , Cl^-) transport. Although a diuretic effect has been described, such an effect is postulated to be secondary to direct renovascular effects (elevated GFR)
- 30 • metabolism:
urotensin-II influences prolactin secretion and exhibits a lipolytic effect in fish (activating triacylglycerol lipase resulting in the mobilization of non-esterified free fatty acids)
(Pearson, *et. al. Proc. Natl. Acad. Sci. (U.S.A.)* 1980, 77, 5021; Conlon, *et. al. J. Exp. Zool.* 1996, 275, 226.)

In studies with human Urotensin-II it was found that it:

- 35 • was an extremely potent and efficacious vasoconstrictor

- exhibited sustained contractile activity that was extremely resistant to wash out
- had detrimental effects on cardiac performance (myocardial contractility)

Human Urotensin-II was assessed for contractile activity in the rat-isolated aorta and was shown to be the most potent contractile agonist identified to date. Based on the *in vitro*

5 pharmacology and *in vivo* hemodynamic profile of human Urotensin-II it plays a pathological role in cardiovascular diseases characterized by excessive or abnormal vasoconstriction and myocardial dysfunction. (Ames *et. al. Nature* 1999, 401, 282)

Compounds that antagonize the Urotensin-II receptor may be useful in the treatment of congestive heart failure, stroke, ischemic heart disease (angina, myocardial
10 ischemia), cardiac arrhythmia, hypertension (essential and pulmonary), COPD, restenosis, asthma, (Hay DWP, Luttmann MA, Douglas SA: 2000, Br J Pharmacol: In press.) neurogenic inflammation and metabolic vasculopathies all of which are characterized by abnormal vasoconstriction and/or myocardial dysfunction. Since U-II and GPR14 are both expressed within the mammalian CNS (Ames *et. al. Nature* 1999, 401, 282), they also may be useful in the
15 treatment of addiction, schizophrenia, impulsivity, anxiety, stress, depression, and neuromuscular function. Functional U-II receptors are expressed in rhabdomyosarcomas cell lines and therefore may have oncological indications. Urotensin may also be implicated in various metabolic diseases such as diabetes (Ames *et. al. Nature* 1999, 401, 282, Nothacker et al., *Nature Cell Biology* 1: 383-385, 1999).

20

SUMMARY OF THE INVENTION

In one aspect this invention provides for quinolines and pharmaceutical compositions containing them.

In a second aspect, this invention provides for the use of quinolines as antagonists of
25 urotensin II, and as inhibitors of urotensin II.

In another aspect, this invention provides for the use of quinolines for treating conditions associated with urotensin II imbalance.


In an yet another aspect, this invention provides for the use of these quinolones analogs for the treatment of congestive heart failure, stroke, ischemic heart disease (angina, myocardial
30 ischemia), cardiac arrhythmia, hypertension (essential and pulmonary), COPD, restenosis, asthma, neurogenic inflammation and metabolic vasculopathies, addiction, schizophrenia, impulsivity, anxiety, stress, depression, neuromuscular function, arthritis and other inflammatory diseases, fibrosis (e.g. pulmonary fibrosis), sepsis, atherosclerosis and dyslipidemiadiabetes, various gastrointestinal dysfunctions such as esophageal reflux and
35 gastric motility disorders.

The urotensin antagonist may be administered alone or in conjunction with one or more other therapeutic agents, said agents being selected from the group consisting of endothelin receptor antagonists, angiotensin converting enzyme (ACE) inhibitors, vasopeptidase inhibitors, diuretics, digoxin, and dual non-selective β -adrenoceptor and α_1 -adrenoceptor antagonists.

Other aspects and advantages of the present invention are described further in the following detailed description of the preferred embodiments thereof.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides for compounds of Formula I:



Formula (I)

wherein:

- R₁ is 1,1-diphenylmethyl, 1,1-diphenylethyl, xanthyl, phenyl, benzimidazolyl, thiophenyl, 3-indolyl, or 2-indolyl, all of which may be substituted or unsubstituted by one, two, or three halogen, C₁₋₆ alkoxy, C₁₋₆ alkyl, benzenesulfonyl, trifluoromethyl, or trifluoromethylthio groups or they may be substituted by a benzyl, which is further substituted or unsubstituted by one, two, or three halogen, C₁₋₆ alkoxy, or C₁₋₆ alkyl groups;
- R₂ is hydrogen or C₁₋₃alkyl;
- R₃ is independently hydrogen, C₁₋₆ alkyl, phenyl, or benzyl, wherein the phenyl or benzyl may be substituted or unsubstituted by a methylenedioxy group, or one or two halogens, C₁₋₃alkyl; or C₁₋₃alkoxy groups; or both R₃ groups together with the carbon they are attached to is a C₃₋₇cycloalkyl group;
- R₄ is hydrogen or C₁₋₃alkyl;
- R₅ is hydrogen, C₁₋₃alkoxy, or CONR₆R₇;
- R₆ is hydrogen or C₁₋₆ alkyl;
- R₇ is hydrogen or C₁₋₆ alkyl; or R₆ and R₇ together with the nitrogen they are attached to form a 5 or 6 membered ring;
- X is -CR₈R₉ or C=O;
- R₈ is hydrogen or C₁₋₃alkyl;
- R₉ is hydrogen or C₁₋₃alkyl;

or R₈ and R₉ together with the carbon they are attached to form a C₅₋₆cycloalkyl group; or a pharmaceutically acceptable salt thereof.

When used herein, the term "alkyl" and similar terms such as "alkoxy" includes all straight chain and branched isomers. Representative examples thereof include methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, *n*-pentyl and *n*-hexyl.

When used herein, the terms 'halogen' and 'halo' include fluorine, chlorine, bromine and iodine and fluoro, chloro, bromo and iodo, respectively.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active form. All of these compounds and their diastereoisomers are contemplated to be within the scope of the present invention.

R₁ is preferably 1,1-diphenylmethyl, xanthyl, phenyl, thiophenyl, 3-indolyl, or 2-indolyl, substituted or unsubstituted by one, two, or three halogen, methoxy, methyl, benzenesulfonyl, trifluoromethyl, or trifluoromethylthio groups, or benzyl, substituted or unsubstituted by one, two, or three halogen groups.

R₂ preferably is hydrogen.

R₃ preferably is hydrogen, C₁₋₃ alkyl, or phenyl or benzyl; or R₃ together with the carbon they are attached to, is a C₅₋₆ cycloalkyl group.

R₄ preferably is hydrogen.

R₅ preferably is methoxy or CONR₆R₇.

R₆ preferably is hydrogen or C₁₋₃ alkyl.

R₇ preferably is hydrogen or C₁₋₃ alkyl.

X preferably is CH₂ or C=O.

Preferred Compounds are:


- N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(4,5-Dibromothiophen-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-*N,N'*-dimethylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(1*H*-Benzoimidazol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N'-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N*-(4-methoxyquinolin-2-yl)-*N*-methylpropane-1,3-diamine;

- (R)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
- (S)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
- 5 *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N,N'*-dimethyl-*N'*-quinolin-2-yl-propane-1,3-diamine;
- N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-quinolin-2-yl-propane-1,3-diamine;
- N*-(1-Benzenesulfonyl-1*H*-indol-3-ylmethyl)-*N'*-quinolin-2-yl-propane-1,3-diamine;
- (R)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
- 10 (S)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
- N*-(1*H*-Indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
- 2-Benzo[1,3]dioxol-5-ylmethyl-*N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
- 15 *N*-(1-Benzenesulfonyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
- 1-Benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;
- 4,6-Dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxyquinolin-2-ylamino)propyl]amide;
- 20 *N*-(4,6-Dichloro-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
- 1-Benzyl-1-*H*-indole-3-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;
- N*-[3-(4-Methoxy-quinolin-2-ylamino)-propyl]-2,2-diphenyl-acetamide;
- N*-(2,2-Diphenyl-ethyl)-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
- N*-[1-(3,5-Dibromobenzyl)-1*H*-indol-3-ylmethyl]-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-
- 25 diamine;
- N*-[2,2-Bis-(4-chlorophenyl)-ethyl]-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
- 2-Benzyl-*N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
- N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-methyl-propane-1,3-
- 30 diamine;
- (1-[[1-Benzyl-1*H*-indol-3-ylmethyl]-amino]-methyl)-cyclohexylmethyl-(4-methoxy-quinolin-2-yl)-amine;
- N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2,2-dimethyl-propane-1,3-diamine;
- 35 2,2-Bis-(4-chlorophenyl)-*N*-[3-(4-methoxy-quinolin-2-ylamino)-propyl]-acetamide;

- 9*H*-Xanthene-9-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;
2-[3-(3,4-Dichloro-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide;
2-[3-[(4,6-Dichloro-1*H*-indol-2-ylmethyl)-amino]-propylamino]-quinoline-4-carboxylic acid dimethylamide;
5 2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide; and
2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid methylamide.
- 10 Most Preferred Compounds are:
N-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide;
2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic
15 acid methylamide;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
N-(1-Benzenesulfonyl-1 *H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
20 *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-methylpropane-1,3-diamine.

Compounds of Formula (I) may be prepared as outlined in the following scheme:

Scheme 1




Conditions: a) Dimethylsulfate, methylene chloride, acetone, reflux; b) trifluoroacetic anhydride, pyridine, rt; c) BocR₂NCH₂CHR₃CH₂NHR₄, acetonitrile, diisopropylethyl amine, reflux; d) 4 N hydrochloric acid in dioxane, rt; e) R₁CHO, acetic acid, sodium methoxide, methanol, rt, then sodium cyanoborohydride. (R₁, R₂, R₃, and R₄ are as defined above.)

Methylation of 2,4-dihydroxyquinoline (1) with dimethylsulfate, followed by treatment with trifluoroacetic anhydride furnished intermediate 2, as outlined in Scheme 1. Coupling of 2 with various mono-*tert*-butoxycarbonyl protected propylenediamines was accomplished in acetonitrile at reflux to give urethanes 3. Removal of the *tert*-butoxycarbonyl protecting group with 4 N hydrochloric acid in dioxane, followed by reductive alkylation of the resultant amines 4 with various aldehydes provided the target compounds 5.

Alternatively, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride mediated coupling of amines 4 with various carboxylic acids gave amides 6, as outlined in Scheme 2. Reduction of the amide carbonyl with borane-tetrahydrofuran also furnished amines 5.


Scheme 2



Conditions: a) $R_1\text{COOH}$, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 1-hydroxybenzotriazole hydrate, *N*-methylmorpholine, *N,N*-dimethylformamide, rt; b) borane in tetrahydrofuran, reflux. (R_1 , R_2 , R_3 , and R_4 are as defined above.)

Compounds of Formula (I) where R_5 is an optionally substituted carboxamide may be prepared as outlined in Scheme 3.

Scheme 3




Conditions: a) PCl_5 , POCl_3 , reflux; b) $R_6R_7\text{NH}$ hydrochloride, triethylamine, chloroform, rt; c) $\text{BocR}_2\text{NCH}_2\text{CHR}_3\text{CH}_2\text{NHR}_4$, triethylamine, ethanol, reflux; d) 4 N hydrochloric acid in dioxane, methylene chloride, rt; e) $R_1\text{CHO}$, acetic acid, methanol, rt, then sodium cyanoborohydride. (R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , and R_7 are as defined above.)

Treatment of hydroxyacid 7 with phosphorus pentachloride and phosphorus oxychloride, followed by reaction with various amines furnished amides 8. Coupling of 8 with various mono-*tert*-butoxycarbonyl protected propylenediamines was accomplished in ethanol at reflux, followed by removal of the *tert*-butoxycarbonyl protecting group with 4 N hydrochloric acid in dioxane to provide amines 9. Reductive alkylation of the resultant amines 9 with various aldehydes provided the target compounds 10.

Compounds of Formula (I) where R₃ = methyl may be prepared as outlined in Scheme

4.
10


Scheme 4



Conditions: a) 3-amino-2-methyl-propionic acid methyl ester, acetonitrile, diisopropylethyl amine, reflux; b) trimethylaluminum, ammonium chloride, methylene chloride, rt; c) borane in tetrahydrofuran, reflux; d) R₁CHO, acetic acid, sodium methoxide, methanol, rt, then sodium cyanoborohydride. (R₁, R₂, R₃, and R₄ are as defined above.)

Coupling of 2 with 3-amino-2-methyl-propionic acid methyl ester (Adams et al. *J. Chem. Soc.*, 1959, 3061) was accomplished in acetonitrile at reflux to give urethane 11. Conversion of ester 11 to amide 12 using trimethylaluminum followed by borane reduction afforded amine 4. Reductive alkylation of amine 4 as described in Scheme 1 (step e) provided the target compound 5.


Compounds of Formula (I) where R₃ = benzyl may be prepared as outlined in Scheme 5.

Scheme 5

Conditions: a) H₂, palladium on carbon, ethanol, hydrochloric acid

- 5 Amide 13 was prepared from (E)-2-cyano-3-phenyl-acrylamide (purchased from Bionet) via hydrogenation. Coupling of 2 with amide 13 was accomplished in acetonitrile at reflux to give urethane 12, which was subsequently converted to the target compound 5 as described in Scheme 4 (steps c and d).

- 10 Compounds of Formula (I) where R₃ = cyclohexyl may be prepared as outlined in Scheme 6.


Scheme 6

Conditions: a) Ammonia, rt; b) hydrogen, palladium on carbon, ethanol, hydrochloric acid

- 15 Conversion of 1-cyano-cyclohexanecarboxylic acid ethyl ester (Julia et al., *Bull. Soc. Chim. Fr.* 1969, 2427) to the corresponding amide using ammonia gas followed by hydrogenation provided amide 15. Coupling of 2 with amide 15 was accomplished in acetonitrile at reflux to give urethane 12, which was subsequently converted to the target compound 5 as described in Scheme 4 (steps c and d).

Scheme 7

Compounds of Formula (I) where R₃ = phenyl may be prepared as outlined in Scheme 7.



5 Conditions: a) Di-tert-butyl dicarbonate, tetrahydrofuran, rt

Mono-protection of 2-phenyl-propane-1,3-diamine (Weinhardt et al. *J. Med. Chem.*, 1985, 28, 694) afforded amine 16. Coupling of 2 with amine 16 was accomplished in acetonitrile at reflux to give urethane 12, which was subsequently converted to the target compound 5 as described in Scheme 4 (steps c and d).

In order to use a compound of the Formula (I) or a pharmaceutically acceptable salt thereof for the treatment of humans and other mammals it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

15 Compounds of Formula (I) and their pharmaceutically acceptable salts may be administered in a standard manner for the treatment of the indicated diseases, for example orally, parenterally, sub-lingually, transdermally, rectally, via inhalation or via buccal administration.

Compounds of Formula (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as syrups, tablets, capsules and lozenges. A syrup formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, peanut oil, olive oil, glycerine or water with a flavoring or coloring agent. Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, terra alba, talc, gelatin, agar, pectin, acacia, stearic acid, starch, lactose and sucrose. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard gelatin capsule shell. Where the composition is in the form of a soft gelatin shell capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of the compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, for example polyethylene glycol, polyvinylpyrrolidone, lecithin, arachis oil, or sesame oil.

- 5 Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered as a dry powder or in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane.

- 10 A typical suppository formulation comprises a compound of Formula (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example polymeric glycols, gelatins, cocoa-butter or other low melting vegetable waxes or fats or their synthetic analogues.

Typical transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

- 15 Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer to themselves a single dose.

- Each dosage unit for oral administration contains suitably from 0.1 mg to 500 mg/Kg, and preferably from 1 mg to 100 mg/Kg, and each dosage unit for parenteral administration contains suitably from 0.1 mg to 100 mg, of a compound of Formula (I) or a pharmaceutically acceptable salt thereof calculated as the free acid. Each dosage unit for intranasal administration contains suitably 1-400 mg and preferably 10 to 200 mg per person. A topical formulation contains suitably 0.01 to 1.0% of a compound of Formula (I).
- 20

- The daily dosage regimen for oral administration is suitably about 0.01 mg/Kg to 40 mg/Kg, of a compound of Formula (I) or a pharmaceutically acceptable salt thereof calculated as the free acid. The daily dosage regimen for parenteral administration is suitably about 0.001 mg/Kg to 40 mg/Kg, of a compound of the Formula (I) or a pharmaceutically acceptable salt thereof calculated as the free acid. The daily dosage regimen for intranasal administration and oral inhalation is suitably about 10 to about 500 mg/person. The active ingredient may be administered from 1 to 6 times a day, sufficient to exhibit the desired activity.
- 25

- 30 These quinoline analogs may be used for the treatment of congestive heart failure, stroke, ischemic heart disease (angina, myocardial ischemia), cardiac arrhythmia, hypertension (essential and pulmonary), COPD, restenosis, asthma, neurogenic inflammation and metabolic vasculopathies, addiction, schizophrenia, impulsivity, anxiety, stress, depression, neuromuscular function, and diabetes.

The urotensin antagonist may be administered alone or in conjunction with one or more other therapeutic agents, said agents being selected from the group consisting of endothelin receptor antagonists, angiotensin converting enzyme (ACE) inhibitors, vasopeptidase inhibitors, diuretics, digoxin, and dual non-selective β -adrenoceptor and α_1 -adrenoceptor antagonists.

No unacceptable toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

The biological activity of the compounds of Formula (I) are demonstrated by the following tests:

Radioligand binding:

HEK-293 cell membranes containing stable cloned human and rat GPR-14 (20 ug/assay) were incubated with 200 pM [125 I] h-U-II (200 Ci/mmol⁻¹ in the presence of increasing concentrations of test compounds in DMSO (0.1 nM to 10 uM), in a final incubation volume of 200 ul (20 mM Tris-HCl, 5 mM MgCl₂). Incubation was done for 30 minutes at room temperature followed by filtration GF/B filters with Brandel cell harvester. 125 I labeled U-II binding was quantitated by gamma counting. Nonspecific binding was defined by 125 I U-II binding in the presence of 100 nM of unlabeled human U-II. Analysis of the data was performed by nonlinear least square fitting.

Ca²⁺-mobilization:

A microtitre plate based Ca²⁺-mobilization FLIPR assay (Molecular Devices, Sunnyvale, CA) was used for the functional identification of the ligand activating HEK-293 cells expressing (stable) recombinant GPR-14. The day following transfection, cells were plated in a poly-D-lysine coated 96 well black/clear plates. After 18-24 hours the media was aspirated and Fluo 3AM-loaded cells were exposed to various concentrations (10 nM to 30 uM) of test compounds followed by h-U-II. After initiation of the assay, fluorescence was read every second for one minute and then every 3 seconds for the following one minute. The inhibitory concentration at 50% (IC₅₀) was calculated for various test compounds.

Inositol phosphates assays:

HEK-293-GPR14 cells in T150 flask were prelabeled overnight with 1 uCi myo-[3 H] inositol per ml of inositol free Dulbecco's modified Eagle's medium. After labeling, the cells were washed twice with Dulbecco's phosphate-buffered saline (DPBS) and then incubated in DPBS containing 10 mM LiCl for 10 min at 37°C. The experiment was initiated by the

addition of increasing concentrations of h-U-II (1 pM to 1 μ M) in the absence and presence of three different concentrations (0.3, 1 and 10 μ M) of test compounds and the incubation continued for an additional 5 min at 37°C after which the reaction was terminated by the addition of 10% (final concentration) trichloroacetic acid and centrifugation. The supernatants were neutralized with 100 μ l of 1M Trizma base and the inositol phosphates were separated on AG 1-X8 columns (0.8 ml packed, 100-200 mesh) in formate phase. Inositol monophosphate was eluted with 8 ml of 200 mM ammonium formate. Combined inositol di and tris phosphate was eluted with 4ml of 1M ammonium formate/ 0.1 M formic acid. Eluted fractions were counted in beta scintillation counter. Based on shift from the control curve K_B was calculated.

- 10 Activity for the compounds of this invention range from (radioligand binding assay):
 $K_i = 1 \text{ nM} - 10000 \text{ nM}$ [e.g. K_i (example 29) = 90 nM]

Example 1

- 15 Preparation of *N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N*-(4-methoxyquinolin-2-yl)propane-1,3-diamine

a) 2-Hydroxy-4-methoxyquinoline

- 20 A slurry of 2,4-dihydroxyquinoline (20.7 g, 0.13mol), potassium carbonate (35.5 g, 0.26mol), and dimethyl sulfate (14.6 ml, 0.15mol) in acetone (800 ml) was heated at reflux for 3 days. The reaction was cooled to ambient temperature then evaporated under reduced pressure. The residue was slurried in a system of water (1000 ml) and ethyl acetate (500 ml) for 1 hour. The solids were collected then rinsed with water (3x250 ml) and ethyl ether (3x250 ml). Vacuum dried over phosphorus pentoxide to give 2-hydroxy-4-methoxyquinoline (16.3 g, 72%) as a tan powder. $[M+H]^+$ 176, $M+CH_3CN=217$.

- 25 b) 1,1,1-Trifluoromethanesulfonic acid 4-methoxyquinolin-2-yl ester

- 30 A slurry of 2-hydroxy-4-methoxyquinoline (13.4 g, 76.6 mmol) in pyridine (75 ml) was slowly treated under argon with trifluoromethanesulfonic anhydride (15.5 ml, 91.2 mmol). The reaction was allowed to stir at ambient temperature. After 4 days, the reaction was evaporated under reduced pressure to an oil that was azeotroped from toluene (2x200 ml) to give the crude product as a brown solid. Flash chromatography on silica (1:1 ethyl acetate/hexanes as eluent) gave 1,1,1-trifluoromethanesulfonic acid 4-methoxyquinolin-2-yl ester (20.4 g, 87%) as a yellow oil that solidified on standing. $[M+H]^+$ 308, $M+CH_3CN=349$.

- 35 c) [3-(4-Methoxyquinolin-2-ylamino)propyl]carbamic acid tert-butyl ester

A solution of 1,1,1-trifluoromethanesulfonic acid 4-methoxyquinolin-2-yl ester (4.51 g, 14.7 mmol), tert-butyl *N*-(3-aminopropyl)carbamate (3.07 g, 17.6 mmol), and diisopropylethylamine (3.84 ml, 22.0 mmol) in anhydrous acetonitrile (35 ml) was heated at reflux for 6 days. The solution was cooled to ambient temperature then evaporated under reduced pressure to an oil. It was taken into water (35 ml) then extracted into ethyl acetate. The extracts were dried (sodium sulfate) then concentrated to an oil. Column chromatography on silica (1:1 ethyl acetate/hexanes) gave [3-(4-methoxyquinolin-2-ylamino)propyl]carbamic acid tert-butyl ester (3.10 g, 64%) as a colorless oil that solidified on standing. $[M+H]^+$ 332.

10 d) *N'*-(4-Methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride

A solution of [3-(4-methoxyquinolin-2-ylamino)propyl]carbamic acid tert-butyl ester (3.10 g, 9.35 mmol) in anhydrous dichloromethane (20 ml) was treated with 4N HCl in 1,4-dioxane (10 ml). It was stirred at ambient temperature for 2 hours then was evaporated under reduced pressure to give *N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride (2.81 g, 99%) as a white solid. $[M+H]^+$ 232.

e) *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine

A solution of *N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride (200 mg, 0.66 mmol) [4a] in methanol (50 ml) was treated with glacial acetic acid (20 drops) and sodium methoxide (95%, 71 mg, 1.31 mmol) followed by a solution of 1-benzyl-3-indole carboxaldehyde (155 mg, 0.66 mmol) in methanol (5.0 ml). The reaction stirred at ambient temperature for 24 hours then was treated with a solution of sodium cyanoborohydride (83 mg, 1.31 mmol) in methanol (2.0 ml). The reaction stirred at ambient temperature for 24 hours. The solution was evaporated under reduced pressure to a residue that was taken into a mixture of saturated aqueous sodium chloride (20 ml) and 10% sodium hydroxide solution (20 ml). It was extracted in ethyl acetate and the extracts were dried (sodium sulfate) then concentrated to an oil. Column chromatography on silica (95/5 dichloromethane/methanolic ammonia) gave *N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine (126 mg, 84%) as a white solid. $[M+H]^+$ 451.

Compounds derived from Scheme 1:

Example	Compound	MS (ES+) m/e [M+H] ⁺
	<i>N</i> -(4,5-Dibromothiophen-2-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-	

2	2-yl)propane-1,3-diamine	486
3	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine	493
4	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)- <i>N,N'</i> -dimethylpropane-1,3-diamine	479
5	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine	527
6	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-2-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	451
7	<i>N</i> -(1 <i>H</i> -Benzoimidazol-2-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	362
8	<i>N'</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N</i> -(4-methoxyquinolin-2-yl)- <i>N</i> -methylpropane-1,3-diamine	465
9	(<i>R</i>)- <i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine	493
10	(<i>S</i>)- <i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine	493
11	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N,N'</i> -dimethyl- <i>N'</i> -quinolin-2-yl-propane-1,3-diamine	449
12	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -quinolin-2-yl-propane-1,3-diamine	421
13	<i>N</i> -(1-Benzenesulfonyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -quinolin-2-yl-propane-1,3-diamine	471
14	(<i>R</i>)- <i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine	527
15	(<i>S</i>)- <i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine	527
16	<i>N</i> -(1 <i>H</i> -Indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	361
17	2-Benzo[1,3]dioxol-5-ylmethyl- <i>N</i> -(1-benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	585
18	<i>N</i> -(1-Benzenesulfonyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-propane-1,3-diamine	501

19	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)-2-methylpropane-1,3-diamine	465
20	(1-{[(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)amino]methyl}cyclohexylmethyl)-(4-methoxyquinolin-2-yl)amine	519
21	<i>N</i> -[1-(3,5-Dibromobenzyl)-1 <i>H</i> -indol-3-ylmethyl]- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	609
22	<i>N</i> -(1-Benzyl-1 <i>H</i> -indol-3-ylmethyl)- <i>N'</i> -(4-methoxy-quinolin-2-yl)-2,2-dimethyl-propane-1,3-diamine	479

Example 23

Preparation of 1-Benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide.

5

A solution of *N*¹-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride (193 mg, 0.63 mmol) [from example 1d], 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid (203 mg, 0.63 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (182 mg, 0.95 mmol), 1-hydroxybenzotriazole hydrate (94 mg, 0.70 mmol), and *N*-methymorpholine (244μl, 2.22 mmol) in anhydrous *N,N*-dimethylformamide (5 ml) was stirred for 2 days. The resulting slurry was taken into water (50 ml) and 10% sodium hydroxide solution (10 ml). Extracted into ethyl acetate. The extracts were dried (sodium sulfate) and evaporated to an oil. Column chromatography on silica (2:1 ethyl acetate/hexanes) gave 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide (269 mg, 80%) as a white solid. [M+H]⁺ 533.

15

Example 24Synthesis of *N*-(1-Benzyl-4,6-dichloro-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride.

- 5 A solution of 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide (200 mg, 0.37 mmol) [from example 12] in anhydrous tetrahydrofuran (15 ml) was treated with 1 *N* borane-tetrahydrofuran complex (1.1 ml, 1.1 mmol). The solution was heated a reflux for 24 hours then cooled to ambient temperature. Treated with the dropwise addition of concentrated hydrochloric acid (1 ml) then stirred for 1
- 10 hour. Evaporated under reduced pressure to a residue and taken into a system of saturated aqueous sodium chloride (20 ml) and 10% sodium hydroxide (20 ml). Extracted into ethyl acetate. The extracts were dried (sodium sulfate) then evaporated to an oil. Column chromatography on silica (95/5 methylene chloride/methanolic ammonia) gave *N*-(1-benzyl-4,6-dichloro-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine as a
- 15 colorless resin. This was taken into chloroform (5 ml) then treated with 4*N* hydrogen chloride in 1,4-dioxane (300μl). Evaporation under reduced pressure gave *N*-(1-benzyl-4,6-dichloro-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride (102 mg, 42%) as a white solid. $[M+H]^+$ 519.

Compounds derived from Scheme 2:

Example	Compound	MS (ES+) m/e [M+H] ⁺
25	1-Benzyl-4,6-dichloro-1 <i>H</i> -indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide	533
26	4,6-Dichloro-1 <i>H</i> -indole-2-carboxylic acid [3-(4-methoxyquinolin-2-ylamino)propyl]amide	443
27	<i>N</i> -(4,6-Dichloro-1 <i>H</i> -indol-2-ylmethyl)- <i>N'</i> -(4-methoxyquinolin-2-yl)propane-1,3-diamine	429
28	1-Benzyl-1- <i>H</i> -indole-3-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide	465
29	<i>N</i> -[3-(4-Methoxy-quinolin-2-ylamino)-propyl]-2,2-diphenyl-acetamide	426
30	<i>N</i> -(2,2-Diphenyl-ethyl)- <i>N'</i> -(4-methoxy-quinolin-2-yl)-propane-1,3-diamine	412
31	9 <i>H</i> -Xanthene-9-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide	440
32	<i>N</i> -[2,2-Bis-(4-chlorophenyl)-ethyl]- <i>N'</i> -(4-methoxy-quinolin-2-yl)-propane-1,3-diamine	480
33	2,2-Bis-(4-chlorophenyl)- <i>N</i> -[3-(4-methoxy-quinolin-2-ylamino)-propyl]-acetamide	494

Example 34

5 Synthesis of 2-[3-(4-Chloro-3-trifluoromethylbenzylamino)propylamino]quinoline-4-carboxylic acid dimethylamide.

a) 2-Chloroquinoline-4-carboxylic acid dimethylamide

10 A slurry of 2-hydroxyquinoline-4-carboxylic acid (3.0 g, 15.9 mmol), phosphorus pentachloride (3.5 g, 16.7 mmol), and phosphorus oxychloride (20 ml) was heated at reflux for 18 hours. The reaction was then cooled to ambient temperature and evaporated under reduced pressure to a black tar. Residue was dissolved in diethyl ether (100 ml) and washed with ice-cold water and ice-cold brine. Dried over anhydrous sodium sulfate and activated charcoal. Filtration through Celite, followed by evaporation under reduced pressure gave 2-chloroquinoline-4-carbonyl chloride as a light green-gray powder.

A solution of the resultant 2-chloroquinoline-4-carbonyl chloride in chloroform (50 ml) was treated with triethylamine (4.4 ml, 31.5 mmol), followed by solid dimethylamine hydrochloride (1.28 g, 15.7 mmol). The reaction stirred for one hour at ambient temperature at which time it was washed with 10% aqueous sodium hydroxide and 15% aqueous citric acid.

- 5 Drying over anhydrous sodium sulfate, followed by evaporation under reduced pressure gave 2-chloroquinoline-4-carboxylic acid dimethylamide (1.5 g, 40%) as a tan solid. $[M+H]^+$ 235.

b) 2-(3-Aminopropylamino)quinoline-4-carboxylic acid dimethylamide dihydrochloride

- 10 A solution of 2-chloroquinoline-4-carboxylic acid dimethylamide (512 mg, 2.18 mmol), tert-butyl N-(3-aminopropyl)carbamate (571 μ l, 3.27 mmol), and triethylamine (608 μ l, 4.36 mmol) in absolute ethanol (35 ml) was heated at reflux for 7 days. The reaction was then allowed to cool to ambient temperature and was evaporated under reduced pressure to give a yellow oil. The residue was purified by column chromatography (ethyl acetate), dissolved in methylene chloride (10 mL), and treated with 4 N hydrogen chloride in dioxane (3 mL). The reaction was maintained at ambient temperature for 24 hours. Evaporation under reduced pressure gave 2-(3-aminopropylamino)quinoline-4-carboxylic acid dimethylamide dihydrochloride (218 mg, 30%) as a pale yellow solid. $[M+H]^+$ 273.

- 20 c) 2-[3-(4-Chloro-3-trifluoromethylbenzylamino)propylamino]quinoline-4-carboxylic acid dimethylamide dihydrochloride

- Following the procedure of 1e, except substituting 2-(3-aminopropylamino)quinoline-4-carboxylic acid dimethylamide dihydrochloride (50 mg, 0.14 mmol) for *N*¹-(4-methoxyquinolin-2-yl)propane-1,3-diamine dihydrochloride, 2-[3-(4-Chloro-3-trifluoromethylbenzylamino)propylamino]quinoline-4-carboxylic acid dimethylamide (35 mg, 47%) was obtained as a colorless oil. $[M+H]^+$ 465.

Example	Compound	MS (ES+) m/e $[M+H]^+$
35	2-[3-(3,4-Dichloro-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide	431
36	2-[3-[(4,6-Dichloro-1 <i>H</i> -indol-2-yl)methyl]-amino]-propylamino]-quinoline-4-carboxylic acid dimethylamide	471
37	2-[3-(4-Chloro-3-trifluoromethylbenzyl-amino)-propylamino]quinoline-4-carboxylic acid methylamide	451

Example 38Synthesis of *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-methyl-propane-1,3-diamine

5 a) 3-(4-Methoxy-quinolin-2-ylamino)-2-methyl-propionic acid methyl ester

A solution of 1,1,1-trifluoromethanesulfonic acid 4-methoxyquinolin-2-yl ester [from Example 1b] (2.03 g, 6.61 mmol) and 3-amino-2-methyl-propionic acid methyl ester (1.5 eq, 1.52 g, 9.91 mmol) in acetonitrile (25 mL) was heated at reflux for five days. It was cooled to room temperature and evaporated to an oil. The crude product was purified via column

10 chromatography on silica (1:1 ethyl acetate/hexane) to give the product (0.75 g, 42%) as an orange oil. $[M+H]^+$ 275.

b) 3-(4-Methoxy-quinolin-2-ylamino)-2-methyl-propionamide

A slurry of ammonium chloride (0.38 g, 7.11 mmol) in dry methylene chloride (20 mL) at 0 °C was treated with trimethylaluminum (1 eq, 7.11 mmol, 3.6 mL of a 2M solution in toluene). The mixture was allowed to warm to room temperature over a period of two hours, at which time a solution of 3-(4-methoxy-quinolin-2-ylamino)-2-methyl-propionic acid methyl ester (0.33 eq, 2.37 mmol, 0.65 g) in dry methylene chloride (5 mL) was added. The solution stirred at room temperature for two hours, then was quenched slowly by addition of

20 concentrated HCl (1 mL). After stirring for an additional 30 minutes, the mixture was diluted with 10% NaOH (30 mL) and brine (20 mL) and extracted into methylene chloride. The organic extracts were dried over Na₂SO₄, filtered and concentrated to an oil. Purification via column chromatography on silica (95:5 methylene chloride/methanolic ammonia) provided the product (0.29 g, 48%) as a white solid. $[M+H]^+$ 260.

25

c) *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-methyl-propane-1,3-diamine

Following the procedures of Example 24 and Example 1e, except substituting 3-(4-methoxy-quinolin-2-ylamino)-2-methyl-propionamide for 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide, the title compound was

30 prepared as a white solid (0.23, 49%). $[M+H]^+$ 465.

Example 39Synthesis of *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-benzyl-propane-1,3-diamine

5 a) 3-Amino-2-benzyl-propionamide

A solution of (E)-2-cyano-3-phenyl-acrylamide (0.50 g, 2.91 mmol) was dissolved in ethanol (25 mL). Concentrated HCl (1 mL) was added, followed by 10% degussa palladium on carbon (1.0 g). The mixture was subjected to hydrogenation conditions (55psi) for 24 hours. It was filtered through Celite and concentrated to give the product (0.62 g, 100%) as a white
10 solid. $[M+H]^+$ 179.

b) 2-Benzyl-3-(4-methoxy-quinolin-2-ylamino)-propionamide

Following the procedure of Example 1c, except substituting 3-amino-2-benzyl-propionamide for tert-butyl *N*-(3-aminopropyl)carbamate, the product (0.13 g, 28%) was
15 obtained as a white solid. $[M+H]^+$ 336.

c) 2-Benzyl-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine

Following the procedure of Example 24, except substituting 2-benzyl-3-(4-methoxy-quinolin-2-ylamino)-propionamide for 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide, the product (0.12 g, 100%) was obtained as a
20 colorless oil. $[M+H]^+$ 322.

d) *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-benzyl-propane-1,3-diamine

Following the procedure of Example 1e, except substituting 2-benzyl-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine for *N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine, the
25 title compound (0.03 g, 15%) was obtained as a white solid. $[M+H]^+$ 541.

Example 4030 Synthesis of (1-[(1-Benzyl-1*H*-indol-3-ylmethyl)-amino]methyl)-cyclohexylmethyl)-(4-methoxy-quinolin-2-yl)-amine

a) 1-Cyano-cyclohexanecarboxylic acid amide

1-Cyano-cyclohexanecarboxylic acid ethyl ester (11.4 g, 63.0 mmol) was placed into a
35 glass pressure vessel and cooled to -78 °C. It was charged with ammonia via needle until the

total volume had doubled. The vessel was sealed and allowed to stir at room temperature for 20 hours. The solvent was evaporated and the resulting slurry triturated with ethyl acetate. The solids were collected by filtration and then filtered through a silica funnel (50g silica), washing with ethyl acetate. The filtrate was concentrated to give the product (2.51 g, 26%) as a white solid. $[M+H]^+$ 153.

b) 1-Aminomethyl-cyclohexanecarboxylic acid amide hydrochloride

A solution of 1-cyano-cyclohexanecarboxylic acid amide (2.34 g, 20.0 mmol) in ethanol (50 mL) was treated with concentrated HCl (3 mL) and 10% degussa palladium on carbon (0.50 g). The mixture was subjected to hydrogenation conditions (50 psi) for 24 hours, then filtered through Celite. The filtrate was concentrated, then azeotroped with methanol to give a viscous oil. The oil was then resubjected to the above hydrogenation conditions for an additional 20 hours. It was filtered and concentrated as above to give the product (3.85 g, 100%) as a white solid. $[M+H]^+$ 157.

c) 1-[(4-Methoxy-quinolin-2-ylamino)-methyl]-cyclohexanecarboxylic acid amide

Following the procedure of Example 1c, except substituting 1-aminomethyl-cyclohexanecarboxylic acid amide hydrochloride for tert-butyl *N*-(3-aminopropyl)carbamate, the product (2.11 g, 59%) was obtained as a white solid. $[M+H]^+$ 314.

d) (1-Aminomethyl-cyclohexylmethyl)-(4-methoxy-quinolin-2-yl)-amine

Following the procedure of Example 24, except substituting 1-[(4-methoxy-quinolin-2-ylamino)-methyl]-cyclohexanecarboxylic acid amide for 1-benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide, the product (2.10 g, 97%) was obtained as a yellow solid. $[M+H]^+$ 300.

e) (1-[(1-Benzyl-1*H*-indol-3-ylmethyl)-amino]methyl)-cyclohexylmethyl)-(4-methoxy-quinolin-2-yl)-amine

Following the procedure of Example 1e, except substituting (1-aminomethyl-cyclohexylmethyl)-(4-methoxy-quinolin-2-yl)-amine for *N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine, the title compound (0.05 g, 20%) was obtained as a pale yellow powder. $[M+H]^+$ 519.

Example 41Synthesis of *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-phenyl-propane-1,3-diamine

- 5 a) (3-Amino-2-phenyl-propyl)-carbamic acid tert-butyl ester

A solution of 2-phenyl-propane-1,3-diamine (2.20 g, 14.7 mmol) in dry tetrahydrofuran (70 mL) was cooled to 0 °C and treated over 30 minutes with a solution of di-tert-butyl dicarbonate (0.33 eq, 4.88 mmol, 1.10 g) in dry tetrahydrofuran (20 mL). The mixture was allowed to warm to room temperature, stirring overnight. The thick slurry was then
10 concentrated to a white residue, taken into water, and extracted into ethyl acetate. The organic extracts were washed with brine (2x), dried over Na₂SO₄, filtered, and concentrated to a colorless oil (1.16 g, 95%). [M+H]⁺ 251.

- 15 b) [3-(4-Methoxy-quinolin-2-ylamino)-2-phenyl-propyl]-carbamic acid tert-butyl ester

Following the procedure of Example 1c, except substituting (3-amino-2-phenyl-propyl)-carbamic acid tert-butyl ester for tert-butyl *N*-(3-aminopropyl)carbamate, the product (0.54 g, 38%) was obtained as a white foamy solid. [M+H]⁺ 408.

- 20 c) *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-2-phenyl-propane-1,3-diamine

Following the procedures of Example 1d and 1e, except substituting [3-(4-methoxy-quinolin-2-ylamino)-2-phenyl-propyl]-carbamic acid tert-butyl ester for *N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine, the title compound (0.15 g, 60%) was obtained as a white solid. [M+H]⁺ 527.

25

Example 42

Formulations for pharmaceutical use incorporating compounds of the present invention can be prepared in various forms and with numerous excipients. Examples of such formulations are given below.

30

<u>Tablets/Ingredients</u>	<u>Per Tablet</u>
1. Active ingredient (Cpd of Form. I)	40 mg
2. Corn Starch	20 mg
5 3. Alginic acid	20 mg
4. Sodium Alginate	20 mg
5. Mg stearate	<u>1.3 mg</u>
	2.3 mg

10 Procedure for tablets:

Step 1 Blend ingredients No. 1, No. 2, No. 3 and No. 4 in a suitable mixer/blender.

Step 2 Add sufficient water portion-wise to the blend from Step 1 with careful mixing after each addition. Such additions of water and mixing until the mass is of a consistency to permit its conversion to wet granules.

15 Step 3 The wet mass is converted to granules by passing it through an oscillating granulator using a No. 8 mesh (2.38 mm) screen.

Step 4 The wet granules are then dried in an oven at 140°F (60°C) until dry.

Step 5 The dry granules are lubricated with ingredient No. 5.

Step 6 The lubricated granules are compressed on a suitable tablet press.

20

Inhalant Formulation

A compound of Formula I, (1 mg to 100 mg) is aerosolized from a metered dose inhaler to deliver the desired amount of drug per use.

25 Parenteral Formulation

A pharmaceutical composition for parenteral administration is prepared by dissolving an appropriate amount of a compound of formula I in polyethylene glycol with heating. This solution is then diluted with water for injections Ph Eur. (to 100 ml). The solution is then sterilized by filtration through a 0.22 micron membrane filter and sealed in sterile containers.


30

The above specification and Examples fully disclose how to make and use the compounds of the present invention. However, the present invention is not limited to the particular embodiments described hereinabove, but includes all modifications thereof within the scope of the following claims. The various references to journals, patents and other

publications which are cited herein comprise the state of the art and are incorporated herein by reference as though fully set forth.

What is claimed is:

1. A compound of Formula I:



Formula (I)

5 wherein:

R₁ is 1,1-diphenylmethyl, 1,1-diphenylethyl, xanthyl, phenyl, benzimidazolyl, thiophenyl, 3-indolyl, or 2-indolyl, all of which may be substituted or unsubstituted by one, two, or three halogen, C₁₋₆ alkoxy, C₁₋₆ alkyl, benzenesulfonyl, trifluoromethyl, or trifluoromethylthio groups or they may be substituted by a benzyl, which is further substituted or unsubstituted by one, two, or three halogen,

10 C₁₋₆ alkoxy, or C₁₋₆ alkyl groups;

R₂ is hydrogen or C₁₋₃alkyl;

R₃ is independently hydrogen, C₁₋₆ alkyl, phenyl, or benzyl, wherein the phenyl or benzyl may be substituted or unsubstituted by a methylenedioxy group, or one or two halogens, C₁₋₃alkyl; or C₁₋₃alkoxy groups;

15 or both R₃ groups together with the carbon they are attached to is a C₃₋₇cycloalkyl group;

R₄ is hydrogen or C₁₋₃alkyl;

R₅ is hydrogen, C₁₋₃alkoxy, or CONR₆R₇;

R₆ is hydrogen or C₁₋₆ alkyl;

R₇ is hydrogen or C₁₋₆ alkyl;

20 or R₆ and R₇ together with the nitrogen they are attached to form a 5 or 6 membered ring;

X is -CR₈R₉ or C=O;

R₈ is hydrogen or C₁₋₃alkyl;

R₉ is hydrogen or C₁₋₃alkyl;

or R₈ and R₉ together with the carbon they are attached to form a C₅₋₆cycloalkyl group;

25 or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1 wherein R₁ is 1,1-diphenylmethyl, xanthyl, phenyl, thiophenyl, 3-indolyl, or 2-indolyl, substituted or unsubstituted by one, two, or three halogen, methoxy, methyl, benzenesulfonyl, trifluoromethyl, or trifluoromethylthio groups, or benzyl, substituted or unsubstituted by one, two, or three halogen groups; R₂ is hydrogen; R₃ is hydrogen, C₁₋₃ alkyl, or phenyl or benzyl; or R₃ together with the carbon they are attached to,

is a C₅₋₆ cycloalkyl group; R₄ is hydrogen; R₅ is methoxy or CONR₆R₇; R₆ is hydrogen or C₁₋₃ alkyl; R₇ is hydrogen or C₁₋₃ alkyl; and X is CH₂ or C=O.

3. A compound of claim 1 chosen from the group consisting of:

- 5 *N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(4,5-Dibromothiophen-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-*N,N'*-dimethylpropane-1,3-diamine;
- 10 *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(1*H*-Benzoimidazol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N'-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N*-(4-methoxyquinolin-2-yl)-*N*-methylpropane-1,3-diamine;
- 15 (R)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
(S)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
N-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N,N'*-dimethyl-*N'*-quinolin-2-yl-propane-1,3-diamine;
- 20 *N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-quinolin-2-yl-propane-1,3-diamine;
N-(1-Benzenesulfonyl-1*H*-indol-3-ylmethyl)-*N'*-quinolin-2-yl-propane-1,3-diamine;
(R)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
(S)-*N*-(1-Benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
- 25 *N*-(1*H*-Indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
2-Benzo[1,3]dioxol-5-ylmethyl-*N*-(1-benzyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
N-(1-Benzenesulfonyl-1*H*-indol-3-ylmethyl)-*N'*-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
- 30 1-Benzyl-4,6-dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;
4,6-Dichloro-1*H*-indole-2-carboxylic acid [3-(4-methoxyquinolin-2-ylamino)propyl]amide;
N-(4,6-Dichloro-1*H*-indol-2-ylmethyl)-*N'*-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
- 35 1-Benzyl-1-*H*-indole-3-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;

- N-[3-(4-Methoxy-quinolin-2-ylamino)-propyl]-2,2-diphenyl-acetamide;
 N-(2,2-Diphenyl-ethyl)-N'-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
 N-[1-(3,5-Dibromobenzyl)-1*H*-indol-3-ylmethyl]-N'-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
 5 N-[2,2-Bis-(4-chlorophenyl)-ethyl]-N'-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
 2-Benzyl-N-(1-benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
 N-(1-Benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxy-quinolin-2-yl)-2-methyl-propane-1,3-diamine;
 10 (1-[(1-Benzyl-1*H*-indol-3-ylmethyl)-amino]-methyl)-cyclohexylmethyl)-(4-methoxy-quinolin-2-yl)-amine;
 N-(1-Benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxy-quinolin-2-yl)-2,2-dimethyl-propane-1,3-diamine;
 2,2-Bis-(4-chlorophenyl)-N-[3-(4-methoxy-quinolin-2-ylamino)-propyl]-acetamide;
 15 9*H*-Xanthene-9-carboxylic acid [3-(4-methoxy-quinolin-2-ylamino)-propyl]-amide;
 2-[3-(3,4-Dichloro-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide;
 2-[3-[(4,6-Dichloro-1*H*-indol-2-ylmethyl)-amino]-propylamino]-quinoline-4-carboxylic acid dimethylamide;
 2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid
 20 dimethylamide; and
 2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid methylamide.

4. A compound of Claim 3 chosen from the group consisting of:

- 25 N-(1-benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxyquinolin-2-yl)propane-1,3-diamine;
 2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid dimethylamide;
 2-[3-(4-Chloro-3-trifluoromethyl-benzylamino)-propylamino]-quinoline-4-carboxylic acid methylamide;
 30 N-(1-Benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxyquinolin-2-yl)-2-propylpropane-1,3-diamine;
 N-(1-Benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxyquinolin-2-yl)-2-phenylpropane-1,3-diamine;
 N-(1-Benzenesulfonyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxy-quinolin-2-yl)-propane-1,3-diamine;
 N-(1-Benzyl-1*H*-indol-3-ylmethyl)-N'-(4-methoxyquinolin-2-yl)-2-methylpropane-1,3-diamine.

5. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

6. A method for treating conditions associated with Human Urotensin II imbalance by administering to a subject in need thereof an effective amount of a compound of claim 1.

- 5 7. A method for treating congestive heart failure, stroke, ischemic heart disease (angina, myocardial ischemia), cardiac arrhythmia, hypertension (essential and pulmonary), COPD, restenosis, asthma, neurogenic inflammation and metabolic vasculopathies, addiction, schizophrenia, impulsivity, anxiety, stress, depression, neuromuscular function, and diabetes by administering to a subject in need thereof an effective amount of a compound of claim 1.

10

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/02007

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/47, 31/4709; C07D 215/38, 401/12, 405/12, 409/12; A61P 9/00
 US CL : 514/313; 546/153, 159, 163

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 U.S. : 514/313; 546/153, 159, 163

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,093,333 A (SAAB) 03 March 1992 (03.03.92), column 2.	1-5
A	WO 99/55677 A1 (SMITHKLINE BEECHAM PLC) 04 November 1999 (04.11.99), pages 1-2.	1-5
A	US 6,075,137 A (CULP et al) 13 June 2000 (13.06.00), column 1.	6, 7

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

16 April 2002 (16.04.2002)

Date of mailing of the international search report

11 JUN 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Evelyn Huang *Telicia D. Roberts for*

Telephone No. 703-308-1235

Form PCT/ISA/210 (second sheet) (July 1998)

Best Available Copy